

Your choices, your way with the Chiesi CareDirect® Wellness Program

CHIESI
CAREDIRECT.
WELLNESS

Nutritional Program*

Earn nutritional supplements, probiotics, and vitamins with every approved prescription.



OR



Wellness Points Program*

Redeem points for items such as medical equipment, masks, smart scales, and fitness equipment.


For eligible patients, enrolling is as easy as...

1



Upload a prescription proof of purchase for eligibility and access

2



Choose the Nutritional Program or Wellness Points Program

3



Get vitamins and supplements OR earn points for wellness items

Get the most out of the program with both prescriptions

Pertzye®
(pancrelipase)
Delayed-Release Capsules



Bronchitol®
(mannitol) inhalation powder



Scan to learn more about the savings and support program

Simple & convenient mobile app for patients taking PERTZYE® (pancrelipase)[†]



Scan to download the app and enroll today

Enroll in the Wellness Program

Upload monthly prescriptions

Track your chosen items and balances

*Restrictions apply. Patients receiving Medicare, Medicaid, or that are participating in any other state or federally subsidized pharmacy benefit program are not eligible for this program. Please see full Terms and Conditions for additional eligibility requirements.

†Restrictions apply. Patients receiving Medicare, Medicaid, or that are participating in any other state or federally subsidized pharmacy benefit program are not eligible for the Wellness Program. Please see full Terms and Conditions for additional eligibility requirements and information.

Important Safety Information

The most common side effects of PERTZYE® (pancrelipase) include diarrhea, upset stomach (indigestion), and cough.

The most common side effects of BRONCHITOL® (mannitol) inhalation powder include, cough, coughing up of blood, pain or irritation in the back of your mouth and throat and discomfort when swallowing, vomiting, fever, joint pain, bacteria in your sputum.

Please see Important Safety Information on back.

See Terms and Conditions in pocket.



Uses for PERTZYE[®] (pancrelipase) Delayed-Release Capsules

PERTZYE is a prescription medicine used to treat people who can't digest food normally because their pancreas does not make enough enzymes due to cystic fibrosis or other conditions.

PERTZYE capsules contain a mixture of digestive enzymes, including lipases, proteases, and amylases, from pig pancreas. PERTZYE is safe and effective in children and adults when taken as prescribed by your doctor.

Important Safety Information

PERTZYE may increase the risk of having a rare bowel disorder called fibrosing colonopathy, especially if taken at a high dose in children with cystic fibrosis who are less than 12 years of age. This condition is serious and may require surgery. The risk of having fibrosing colonopathy may be reduced by following the dosing instructions that your doctor gives you. Call your doctor right away if you have any unusual or severe stomach area (abdominal) pain, bloating, trouble passing stools (constipation), nausea, vomiting, or diarrhea.

Take PERTZYE capsules exactly as your doctor tells you. You should not switch PERTZYE with any other pancreatic enzyme product without first talking with your doctor. Don't take more capsules in a day than the number your doctor tells you to take (total daily dose). Always take PERTZYE with a meal or snack and plenty of fluid. If you eat a lot of meals or snacks in a day, be careful not to go over your total daily enzyme dose. PERTZYE capsules should be swallowed whole. Do not crush or chew the PERTZYE capsules or their contents, and do not hold the capsule or capsule contents in your mouth. Read the Medication Guide for instructions on how to take PERTZYE if you have trouble swallowing capsules.

The most common side effects of PERTZYE include diarrhea, upset stomach (indigestion), and cough. Other potential serious side effects are irritation of the inside of your mouth; increase in blood uric acid levels, which may cause worsening of swollen, painful joints (gout); and allergic reactions, including trouble swallowing or breathing, skin rash, itching, or swelling of your face, eyes, lips, tongue, or throat. Call your doctor right away if you have any of these symptoms.

Tell your doctor about all of your medical conditions, including if you are allergic to pork (pig) products, have a history of blockage of your intestines, or scarring or thickening of your bowel wall (fibrosing colonopathy), have gout, kidney disease, or high blood uric acid (hyperuricemia), have trouble swallowing capsules, are pregnant or plan to become pregnant, or are breastfeeding or plan to breastfeed. Tell your doctor about all the medicines you take.

PERTZYE and other pancreatic enzyme products are made from the pancreas of pigs, the same pigs people eat as pork. These pigs may carry viruses. Although it has never been reported, it may be possible for a person to get a viral infection from taking pancreatic enzyme products that come from pigs.

Read the **Full Prescribing Information and Medication Guide** before you start taking PERTZYE. You should also read the insert that comes with each prescription refill because there may be new information.

Please see Patient Information within the accompanying Full Prescribing Information.



About BRONCHITOL[®] (mannitol) inhalation powder

BRONCHITOL is a prescription medicine that is used along with other therapies to improve lung function in people 18 years of age and older with cystic fibrosis (CF).

BRONCHITOL is only for adults who have passed the BRONCHITOL Tolerance Test (BTT). Your first dose of BRONCHITOL is given during the BTT by your healthcare provider and tests if BRONCHITOL is right for you. Your healthcare provider will use equipment to monitor you and have medicine ready if you have bronchospasm during the test. If you have bronchospasm during your BTT, then you should not be prescribed BRONCHITOL.

BRONCHITOL should not be used in children and adolescents. It is not known if BRONCHITOL is safe and effective in children under 18 years of age.

Important Safety Information

Do not take BRONCHITOL if you have had an allergic reaction to mannitol or any parts of the BRONCHITOL capsule or if you do not pass the BRONCHITOL Tolerance Test (BTT).

Before you take BRONCHITOL, tell your healthcare provider about all your medical conditions, including if you have ever coughed up blood or had blood in your mucus (sputum); are pregnant or planning on becoming pregnant. It is not known if BRONCHITOL will harm your unborn baby. Tell your healthcare provider right away if you become pregnant while using BRONCHITOL; are breastfeeding or plan to breastfeed. It is not known if BRONCHITOL passes into your breast milk or if it can harm your baby. Talk to your healthcare provider about the best way to feed your baby while using BRONCHITOL.

Tell your healthcare provider about all medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

BRONCHITOL may cause serious side effects, including sudden breathing problems immediately after inhaling your medicine; coughing up of blood (hemoptysis). Call your healthcare provider or get emergency medical care right away if you cough up a large amount of blood.

The most common side effects of BRONCHITOL include, cough, coughing up of blood, pain or irritation in the back of your mouth and throat and discomfort when swallowing, vomiting, fever, joint pain, bacteria in your sputum. Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of BRONCHITOL. You can ask your healthcare provider or pharmacist for more information.

Please see Patient Information within the accompanying Full Prescribing Information.

You are encouraged to report negative side effects of prescription drugs to the FDA.
Visit www.fda.gov/medwatch or call 1-800-FDA-1088.